



ADIMA GLOBAL IMPEX

Poultry Feed Supplement Manufacturer & Importers

Address: Shree Mookambika Complex, First Floor, Pandith Hari Bhat Road, Haleyangadi Post, Mangalore - 574146, Karnataka.

Email: adimaglobalimpex@gmail.com

Mob: Adithya K.N Bharadwaj: 7411468156, 6362137407





Poultry Feed Supplements

Brand

Content

Indication

Dosage

Thylo-Ma

Tylosin Phosphate 10%

Increases Egg Production and Improves Feed Efficiency in Layers & Breeders and for Control of Mycoplasma Infections.

500g to 1kg/Ton of Feed or as Advised by Veterinarian.

Livo-Ma

Livertonic Powder

Protects the Liver from the Effect of Toxins in Poultry. General Livertonic Boosts Liver Function Helps to Overcome Toxicity and Fatty Liver Condition to Minimize & Prevent Liver Damage.

500g Livo-Ma in 1 Ton of Feed.

Coli-Ma

Choline Chloride 60%

To Decrease Chick Mortality During Rapid Growth by Mobilization of Excess Fat from the Liver.

500g to 1kg/Ton of Feed or as Advised by Veterinarian.

ENRA-Ma

Enramycin 8%

For Increase in Rate of Weight Gain. Improved Feed Efficiency and for the Control of Necrotic Enteritis Caused or Complicated by Clostridium spp or other Organisms.

Broiler Starter and Pre-Starter Feed 63 to 125g/Ton of Feed Finisher 38 to 125g/Ton of Feed Layers/Breeders 63 to 125g.

CTC-Ma

Chlortetracycline 15%

To Treat Conjunctivitis and Infected Wounds of Animals. It can Prevent and Cure Disease, Promote Growth with a Specific Effect on Chicken Swine in Poultry.

250g/Per Ton.





Anti - Coccidiosis

Brand

Content

Indication

Dosage

Salino-Ma

Salinomycin 12%

Decrease in Incidence of Necrotis Enteritis in Broiler Chicken. It is mainly used as a Coccidiostat in Poultry to Prevent Coccidiosis and Promote Growth.

500g Salino-Ma in 1 Ton of Feed.

Madu-Ma

Maduramycin 1%

For Use in Broiler Chickens as an Aid in the Prevention of Coccidiosis caused by Eimeria Spicies Etc. It also Enhances Productivity with Higher Safety Margin.

500g/Per Ton. (Make a Pre Mix Before Adding)

Mono-Ma

Monensin Sodium 20%

Intended to be used as a Coccidiostat in Feed for Chickens.

500g/Per Ton of Fineshed Feed.

NARA-Ma 10%

Narasin 10%

Treatment of Coccidiosis and to Prevent Necrotic Enteritis in Broiler Chicken.

500g to 600g of Narsasin 10% in 1 Ton of Feed.

NINA-Ma 8%

Nicarbazine 8% + Narasin 8%

Prevents Coccidiosis and Provides Weight Gain.

500g Nina-Ma in 1 Ton of Feed.



Brand**Content****Indication****Dosage**

Adi-Tim

Tiamulin Hydrogen
Fumarate 10%

Increases Growth Rate
Performance, Feed Efficiency
and Production in Poultry.
To control Mycoplasma and
related Conditions in Poultry.

Tiamulin 10%
-200g/Ton of
Feed and
Administer on
Regular Basis
to Laying Hens.

PhyMa 5000IU

Phytase 5000 IU

This Feeding Especially in the
Poultry Industry to Enhance
Phosphorus Intake and Minimize
Environmental Pollution Derived
from Equalized Source.

100g/Per Ton.

Toxi-Ma

Toxin Binder

Keeps Poultry Feed Healthy
by Absorbing a Wide Range of
Mycotoxins. It is Stable During
Feed Processing and Storage and
has no Binding Affinity for Vitamin,
Minerals, Drugs and Other Nutrients.

500g to 1kg of
Toxi-ma in 1 Ton
of Feed.



Adima Global Impex

GSTIN: 29ABWFA0133HIZC

D.L.No: 20B-KA-MN2-261760 21B-KA-MN2-261761

CONTACT

Email: adimaglobalimpex@gmail.com

Mob: Adithya K.N Bharadwaj
7411468156, 6362137407